

# SOURCE MOLECULAR CORPORATION

4985 SW 74th Court, Miami, FL 33155 USA

Tel: (1) 786-268-8363, Fax: (1) 786-513-2733, Email: info@sourcemolecular.com

## Human Enterococcus ID™

Detection of the *Enterococcus faecium* esp Human Gene Biomarker for Human Fecal Contamination by Polymerase Chain Reaction (PCR) DNA Analytical Technology

Submitter: ABC Beach Park

Submitter #'s: 775, 776, 777 and 778

Source Molecular #'s: SM 0125, SM 0126, SM 0127 and SM 0128

Samples Received: January 3rd, 2011

Date Reported: January 10th, 2011

**SAMPLE**

SM #	Client #	Enterococci (CFU/100mL) <sup>7</sup>	DNA Analytical Results
SM 0125	775	45	Human Gene Biomarker Detected
SM 0126	776	150	Negative
SM 0127	777	255	Human Gene Biomarker Detected
SM 0128	778	15	Negative

### Laboratory Comments

The submitted water samples were filtered for *Enterococcus spp.* and the *Enterococci* were enumerated on petri plates. Afterwards, the *Enterococci* were eluted and centrifuged directly from the filter for DNA analysis.

All reagents, chemicals and apparatuses were verified and inspected beforehand to ensure that no false negatives or positives could be generated. In that regard, positive and negative controls were run to attest the integrity of the analysis. All inspections and controls tested negative for possible extraneous contaminants, including PCR inhibitors.

Samples 776 (Our Ref: SM 0126) and 778 (Our Ref: SM 0128) tested negative for the *Enterococcus faecium* human gene biomarker. It is important to note that a negative result does not mean that the sample does not definitely have human contamination. In order to strengthen the result, a negative sample should be analyzed further for human fecal contamination with other DNA analytical tests such as the Human Bacteroidetes ID™ and Human Fecal Virus ID™ services.

Samples 775 (Our Ref: SM 0125) and 777 (Our Ref: SM 0127) tested positive for the *Enterococcus faecium* human gene biomarker suggesting that human fecal contamination is present in these water samples. The client is nonetheless encouraged to conduct other DNA analytical tests such as the services mentioned above to further confirm the results.

### DNA Analytical Method Explanation

For each sample, 100 ml of water was filtered through a 0.45 micron membrane filter and placed on nEI agar. The samples were incubated for 24 hours. Each filter was removed, placed in a buffer and vortexed vigorously. Once the buffer was spun to pellet the bacteria, the supernatant was removed and the pellet was resuspended in a small volume of water. DNA extraction was prepared using the Qiagen DNA extraction kit, as per manufacturer's instructions.

Five micro-liter aliquots of purified DNA extraction were used directly as template for subsequent PCR reactions. Amplification of PCR primers were carried out using HotStarTaq polymerase (Qiagen, Inc.) and master mix, which contained a final concentration of 1.5 mM MgCl<sub>2</sub>, 150 mM dNTP, and 0.3 mM of each primer.

An Eppendorf Gradient Thermocycler was used with the following cycling parameters: 95°C for 15 minutes (to lyse cells and activate polymerase), followed by 35 cycles of 94°C for 1 minute, 55°C for 1 minute, and 72°C for 1 minute and a final extension at 72°C for 5 minutes. PCR products were electrophoresed on 2% agarose gels, stained with GelStar nucleic acid stain (Cambrex, Inc.) and visualized under UV light.

### DNA Analytical Theory Explanation

*Enterococci* are a subgroup of Fecal *Streptococci* and are characterized by their ability to grow in 6.5% sodium chloride, at low and elevated temperatures (10 °C and 45 °C), and at elevated pH (9.5). These microorganisms have been used as indicators of fecal pollution for many years and have been especially valuable in the marine environment and recreational waters as indicators of potential health risks and swimming-related gastroenteritis.<sup>1</sup>

*Enterococci* are benign bacteria when they reside in their normal habitat such as the gastrointestinal tracts of human or animals. Outside of their normal habitat, *Enterococci* are pathogenic causing urinary tract and wound infections, and life-threatening diseases such as bacteraemia, endocarditis, and meningitis. *Enterococci* easily colonize open wounds and skin ulcers.

Compounding their pathogenesis, *Enterococci* are also some of the most antibiotic resistant bacteria, particularly from human sources. Studies have shown that certain strains of *Enterococci* are resistant to expensive and potent antibiotics such as vancomycin. This is particularly worrisome for the medical community since these antibiotics are given as a last resort to fight severe bacterial infections.

Several intrinsic features of the *Enterococcus* genus allow it to survive for extended periods of time, leading to its extended survivability and diffusion. For example, *Enterococci* have been shown to survive for 30 minutes at 60°C and persist in the presence of detergents. As such, the inherent ruggedness of *Enterococcus* confers it a strong tolerance to many classes of antibiotics.

The Human Enterococcus ID™ service is designed around the principle that certain strains of the *Enterococcus* genus are specific to humans.<sup>2,3,4</sup> These *Enterococci* can be used as indicators of human fecal contamination. Strains of *Enterococcus faecium*, *Enterococcus faecalis* and yellow-pigmented *Enterococci* have been shown to be from human sources.<sup>2,3,4</sup> Within these *Enterococcus spp.* are genes associated with *Enterococci* that are specific to humans.<sup>5</sup> The Human Enterococcus ID™ service targets the esp human gene biomarker in *Enterococcus faecium*.<sup>6</sup>

One of the advantages of the Human Enterococcus ID™ service is that the entire population of *Enterococci* of the selected portion of the water sample is screened. As such, this method avoids the randomness effect of selecting isolates off a petri dish. This is a particular advantage for highly contaminated water systems with potential multiple sources of fecal contamination.

Accuracy of the results is possible because the method uses PCR DNA technology. PCR allows quantities of DNA to be amplified into large number of small copies of DNA sequences. This is accomplished with small pieces of DNA called primers that are complementary and specific to the genomes to be detected.

Through a heating process called thermal cycling, the double stranded DNA is denatured and inserted with complementary primers to create exact copies of the DNA fragment desired. This process is repeated rapidly many times ensuring an exponential progression in the number of copied DNA. If the primers are successful finding a site on the DNA fragment that is specific to the genome to be studied, then billions of copies of the DNA fragment will be available for detection by gel electrophoresis.

The gel electrophoresis apparatus uses an electrical field to distinguish different DNA fragments according to their molecular weights. Lighter DNA fragments will move farther along the gel than their heavier counterparts. At the end of the procedure different bands of accumulated DNA fragments will aggregate at different parts of the gel. It is this accumulation of DNA fragments that creates a band on the gel. Researchers use these bands to distinguish certain genomes such as the human gene biomarker from *Enterococcus faecium*.

These banding patterns confirm or negate the presence of the *Enterococci* human gene biomarker. As such, the banding patterns provide a reliable indicator of human fecal contamination. To strengthen the validity of the results, the Human Enterococcus ID™ service should be combined with other DNA analytical services such as the Human Bacteroidetes ID™ and Human Fecal Virus ID™ services.

<sup>1</sup> Scott, Troy M., Rose, Joan B., Jenkins, Tracie M., Farrah, Samuel R., Lukasik, Jerzy **Microbial Source Tracking: Current Methodology and Future Directions.** Appl. Environ. Microbiol. (2002) 68: 5796-5803.

<sup>2</sup> Wheeler, A.L., P.G. Hartel, D.G. Godfrey, J.L. Hill, and Segars W.I. 2002. **Potential of *Enterococcus faecalis* as a human fecal indicator for microbial source tracking.** J Environ Qual. 31(4):1286-93.

<sup>3</sup> Bahirathan ML, Puente L, Seyfried P. 1998. **Use of yellow-pigmented enterococci as a specific indicator of human and nonhuman sources of faecal pollution.** Can J Microbiol 44:1066-1071.

<sup>4</sup> Quednau, M., Ahrne, S., Molin, G. **Genomic Relationships between *Enterococcus faecium* Strains from Different Sources and with Different Antibiotic Resistance Profiles Evaluated by Restriction Endonuclease Analysis of Total Chromosomal DNA Using EcoRI and PvuII.** Appl. Environ. Microbiol. 1999 65: 1777-1780.

<sup>5</sup> Hammerum, A.M., and L.B. Jensen. 2002. **Prevalence of esp, encoding the enterococcal surface protein, in *Enterococcus faecalis* and *Enterococcus faecium* isolates from hospital patients, poultry, and pigs in Denmark.** J. Clin. Microbiol. 40: 4396.

<sup>6</sup> Scott, T.M., T.M. Jenkins, J. Lukasik, and J.B. Rose. 2005 **Potential Use of a Host Associated Molecular Marker in *Enterococcus faecium* as an Index of Human Fecal Pollution.** Environ. Sci. Technol. 39: 283-287.

<sup>7</sup> EPA Method 1600 (modified): Membrane Filter Test Method for Enterococci In Water (1997).

### Limitation of Damages – Repayment of Service Price

It is agreed that in the event of breach of any warranty or breach of contract, or negligence of the Source Molecular Corporation, as well as its agents or representatives, the liability of the Source Molecular Corporation shall be limited to the repayment, to the purchaser (submitter), of the individual analysis price paid by him/her to the Source Molecular Corporation. The Source Molecular Corporation shall not be liable for any damages, either direct or consequential. The Source Molecular Corporation provides analytical services on a PRIME CONTRACT BASIS ONLY. Terms are available upon request.